

Association Between Arsenic Exposure from a Coal-Burning Power Plant and Urinary Arsenic Concentrations in Prievidza District, Slovakia

Ulrich Ranft, Peter Miskovic, Beate Pesch, Pavel Jakubis, Elenora Fabianova, Tom Keegan, Andre Hergemöller, Marian Jakubis, Mark J Nieuwenhuijsen, and the EXPASCAN Study Group (Bencko V., Colvile R., Cordos E., Docx P., Farago M., Frank P., Götzl M., Grellier J, Hong B., Rames J., Rautiu, R., Stevens E., Thornton I., Unfried K., Zvarova J.)
doi:10.1289/ehp.5838 (available at <http://dx.doi.org/>)
Online 9 January 2003



Association Between Arsenic Exposure from a Coal-Burning Power Plant and Urinary Arsenic Concentrations in Prievidza District, Slovakia

Ulrich Ranft¹, Peter Miskovic², Beate Pesch¹, Pavel Jakubis³, Elenora Fabianova², Tom Keegan⁴, Andre Hergemöller¹, Marian Jakubis³, Mark J Nieuwenhuijsen⁴, and the EXPASCAN Study Group⁵

¹ Institut für Umweltmedizinische Forschung (IUF) an der Heinrich-Heine-Universität Düsseldorf, Germany

² State Health Institute, Banska Bystrica, Slovakia

³ State Health Institute, Prievidza, Slovakia

⁴ Department of Environmental Science and Technology, Imperial College of Science, Technology and Medicine, London, United Kingdom

⁵ Bencko V., Colvile R., Cordos E., Docx P., Farago M., Frank P., Götzl M., Grellier J, Hong B., Rames J., Rautiu, R., Stevens E., Thornton I., Unfried K., Zvarova J.

Reprints and correspondence to

Prof. Dr. Ulrich Ranft

Institut für Umweltmedizinische Forschung (IUF) an der Heinrich-Heine-Universität Düsseldorf, Auf'm Hennekamp 50

40225 Düsseldorf, Germany

phone +49 211 3389 287

fax +49 211 3389 283

ranft@uni-duesseldorf.de

Running title

Environmental and Urinary Arsenic

Key words

Environmental health, arsenic, biomarker, non-melanoma skin carcinoma, environmental exposure, coal combustion

Acknowledgments

Funding was provided by the European Commission under Contract Number IC15 CT98 0525.

List of Abbreviations

As	arsenic
As _{inorg}	inorganic arsenic (As ^{III} +As ^V)
As _{methyl}	methylated arsenic (MMA+DMA)
As _{sum}	sum of speciated arsenic compounds (As _{inorg} +As _{methyl})
CI	confidence interval
Crn	creatinine
DMA	dimethylarsinic acid
EPA	Environmental Protection Agency
EC	European Commission
EXPASCAN	Exposure to arsenic and cancer risks in Central and Eastern Europe
IARC	International Agency for Research on Cancer
LD	Limit of detection

MMA	monomethylarsonic acid
NMSC	non-melanoma skin cancer
R ²	fraction of explained variance
SK NCI	National Cancer Institute of Slovakia

Outline of manuscript section headers

Abstract

Introduction

Material and Methods

Recruitment of the study population

Exposure information

Arsenic speciation in urine samples

Statistical methods

Results

Discussion

Urinary As as biomarker of environmental As exposure

Exposure to environmental arsenic and urinary arsenic excretion

Health effects of arsenic

Conclusions

References

ABSTRACT

To assess the arsenic (As) exposure of a population living in the vicinity of a coal-burning power plant with high arsenic emission in the Prievidza District, Slovakia, 548 spot urine samples were speciated for inorganic As (As_{inorg}), monomethylarsonic and dimethylarsinic acid (MMA, DMA) and their sum (As_{sum}). The urine samples were collected from the study population of a case-control study on non-melanoma skin cancer (NMSC). 411 samples with complete As speciations, sufficient urine quality and without fish consumption were passed on to statistical analysis. Although current environmental As exposure and urinary As concentrations were low (median As in soil 41 $\mu\text{g/g}$ within 5 km distance to the power plant; median urinary As_{sum} 5.8 $\mu\text{g/liter}$), there was a significant but weak association between As in soil and urinary As_{sum} ($r=0.21$, $p<0.01$). A multivariate regression analysis was performed to calculate adjusted regression coefficients for environmental As exposure and other determinants of urinary As. Persons living in the vicinity of the plant had a 27% higher As_{sum} ($p<0.01$), based on elevated concentrations of the methylated species. A 32% increase of MMA occurred among subjects with consumption of home-grown food ($p<0.001$). NMSC cases had significantly higher levels of As_{sum} , DMA and As_{inorg} . The methylation index $As_{inorg}/(MMA+DMA)$ was about 20% lower among cases ($p<0.05$) and in men ($p<0.05$) as compared to controls and females, respectively.